

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use

EXTINA[®] Foam safely and effectively. See full prescribing information for EXTINA[®] Foam.

**EXTINA[®] (ketoconazole) Foam, 2%
For topical use only
Initial U.S. Approval: 1981**

INDICATIONS AND USAGE

EXTINA[®] Foam is indicated for topical treatment of seborrheic dermatitis in immunocompetent patients 12 years of age and older (1).

Safety and efficacy of EXTINA[®] Foam for treatment of fungal infections have not been established.

DOSAGE AND ADMINISTRATION

- EXTINA[®] Foam should be applied to the affected area(s) twice daily for four weeks (2).

- EXTINA[®] Foam is not for ophthalmic, oral, or intravaginal use (2).

DOSAGE FORMS AND STRENGTHS

EXTINA[®] Foam contains 2% ketoconazole in a thermolabile hydroethanolic foam in 50 g and 100 g containers (3).

CONTRAINDICATIONS

None

WARNINGS AND PRECAUTIONS

- EXTINA[®] Foam may result in contact sensitization, including photoallergenicity (5.1, 6.2).
- The contents of EXTINA[®] Foam are flammable (5.2).

ADVERSE REACTIONS

The most common adverse reactions observed in clinical studies (incidence > 1%) were application site burning and application site reaction (6.1).

To report SUSPECTED ADVERSE REACTIONS, contact Stiefel Laboratories, Inc. at 1-888-500-DERM or adverse.event@stiefel.com and FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

To report SUSPECTED ADVERSE REACTIONS, contact at or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling

Revised: 10/2009

FULL PRESCRIBING INFORMATION: CONTENTS *

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Contact Sensitization
- 5.2 Flammable Contents
- 5.3 Systemic Effects

6 ADVERSE REACTIONS

- 6.1 Adverse Reactions in Clinical Trials
- 6.2 Dermal Safety Studies

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use

8.5 Geriatric Use

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics
- 12.4 Microbiology

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

- 17.1 Instructions for Use
- 17.2 Local Reactions

PATIENT INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

EXTINA[®] (ketoconazole) Foam, 2% is indicated for the topical treatment of seborrheic dermatitis in immunocompetent patients 12 years of age and older. Safety and efficacy of EXTINA[®] Foam for treatment of fungal infections have not been established.

2 DOSAGE AND ADMINISTRATION

EXTINA[®] Foam should be applied to the affected area(s) twice daily for four weeks.

Hold the container upright, and dispense EXTINA[®] Foam into the cap of the can or other cool surface in an amount sufficient to cover the affected area(s). Dispensing directly onto hands is not recommended, as the foam will begin to melt immediately upon contact with warm skin. Pick up small amounts of EXTINA[®] Foam with the fingertips, and gently massage into the affected area(s) until the foam disappears. For hair-bearing areas, part the hair, so that EXTINA[®] Foam may be applied directly to the skin (rather than on the hair).

Avoid contact with the eyes and other mucous membranes. EXTINA[®] Foam is not for ophthalmic, oral or intravaginal use.

3 DOSAGE FORMS AND STRENGTHS

EXTINA[®] Foam contains 2% ketoconazole in a thermolabile hydroethanolic foam, and is provided in 50 g and 100 g aluminum containers.

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Contact Sensitization

EXTINA[®] Foam may result in contact sensitization, including photoallergenicity. *[See Adverse Reactions (6.2)]*

5.2 Flammable Contents

The contents of EXTINA[®] Foam include alcohol and propane/butane, which are flammable. Avoid fire, flame and/or smoking during and immediately following application. Do not puncture and/or incinerate the containers. Do not expose containers to heat and/or store at temperatures above 120°F (49°C).

5.3 Systemic Effects

Hepatitis has been seen with orally administered ketoconazole (1:10,000 reported incidence). Lowered testosterone and ACTH-induced corticosteroid serum levels have been seen with high doses of orally administered ketoconazole. These effects have not been seen with topical ketoconazole.

6 ADVERSE REACTIONS

6.1 Adverse Reactions in Clinical Trials

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug, and may not reflect the rates observed in practice. The adverse reaction information from clinical trials does, however, provide a basis for identifying the adverse reactions that appear to be related to drug use and for approximating rates.

The safety data presented in Table 1 (below) reflect exposure to EXTINA[®] Foam in 672 subjects, 12 years and older with seborrheic dermatitis. Subjects applied EXTINA[®] Foam or vehicle foam twice daily for 4 weeks to affected areas on the face, scalp, and/or chest. Adverse reactions occurring in > 1% of subjects are presented in Table 1.

Table 1: Adverse Reactions Reported by > 1% Subjects in Clinical Trials

| Adverse Reactions | EXTINA [®] Foam N = 672 n (%) | Vehicle Foam N = 497 n (%) |
|--|--|----------------------------------|
| Subjects with an Adverse Reaction | 188 (28%) | 122 (25%) |
| Application site burning | 67 (10%) | 49 (10%) |
| Application site reaction | 41 (6%) | 24 (5%) |

Application site reactions that were reported in $\leq 1\%$ of subjects were dryness, erythema, irritation, paresthesia, pruritus, rash and warmth.

6.2 Dermal Safety Studies

In a photoallergenicity study, 9 of 53 subjects (17%) had reactions during the challenge period at both the irradiated and non-irradiated sites treated with EXTINA[®] Foam. EXTINA[®] Foam may cause contact sensitization.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Teratogenic Effects, Pregnancy Category C: Ketoconazole has been shown to be teratogenic (syndactylia and oligodactylia) in the rat when given orally in the diet at 80 mg/kg/day (4.8 times the maximum expected human topical dose based on a mg/m² comparison, assuming 100% absorption from 8 g of foam). However, these effects may be partly related to maternal toxicity, which was also observed at this dose level. [See *Pharmacokinetics (12.3)*]

No reproductive studies in animals have been performed with EXTINA[®] Foam. There are no adequate and well-controlled studies of EXTINA[®] Foam in pregnant women.

EXTINA[®] Foam should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

8.3 Nursing Mothers

It is not known whether EXTINA[®] Foam administered topically could result in sufficient systemic absorption to produce detectable quantities in breast milk. Because many drugs are excreted in human milk, caution should be exercised when EXTINA[®] Foam is administered to women who are breastfeeding.

8.4 Pediatric Use

The safety and effectiveness of EXTINA[®] Foam in pediatric patients less than 12 years of age have not been established.


Of the 672 subjects treated with EXTINA[®] Foam in the clinical trials, 44 (7%) were from 12 to 17 years of age. [See *Clinical Studies (14)*]

8.5 Geriatric Use

Of the 672 subjects treated with EXTINA[®] Foam in the clinical trials, 107 (16%) were 65 years and over.

11 DESCRIPTION

EXTINA[®] Foam contains 2% ketoconazole USP, an antifungal agent, in a thermolabile hydroethanolic foam for topical application. The chemical name for ketoconazole is piperazine, 1-acetyl-4-[4-[[2-(2,4-dichlorophenyl)-2-(1*H*-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]-, *cis*- with the molecular formula C₂₆H₂₈Cl₂N₄O₄ and a molecular weight of 531.43. The following is the chemical structure:



EXTINA[®] Foam contains 20 mg ketoconazole per gram in a thermolabile hydroethanolic foam vehicle consisting of cetyl alcohol, citric acid, ethanol (denatured with *tert*-butyl alcohol and brucine sulfate) 58%, polysorbate 60, potassium citrate, propylene glycol, purified water, and stearyl alcohol pressurized with a hydrocarbon (propane/butane) propellant.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

The mechanism of action of ketoconazole in the treatment of seborrheic dermatitis is not known.

12.2 Pharmacodynamics

The pharmacodynamics of EXTINA[®] Foam has not been established.

12.3 Pharmacokinetics

In a bioavailability study, 12 subjects with moderate to severe seborrheic dermatitis applied 3 g of EXTINA[®] Foam twice daily for 4 weeks. Circulating plasma levels of ketoconazole were < 6 ng/mL for a majority of subjects (75%), with a maximum level of 11 ng/mL observed in one subject.

12.4 Microbiology

Ketoconazole is an antifungal agent which inhibits the *in vitro* synthesis of ergosterol, a key sterol in the cell membrane of *Malassezia furfur*. The clinical significance of antifungal activity in the treatment of seborrheic dermatitis is not known.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies have not been performed to evaluate the carcinogenic or photo-carcinogenic potential of EXTINA[®] Foam. In oral carcinogenicity studies in mice (18-months) and rats (24-months) at dose levels of 5, 20 and 80 mg/kg/day ketoconazole was not carcinogenic. The high dose in these studies was approximately 2.4 to 4.8 times the expected topical dose in humans based on a mg/m² comparison. In a bacterial reverse mutation assay, ketoconazole did not express any mutagenic potential. In three *in vivo* assays (sister chromatid exchange in humans, dominant lethal and micronucleus tests in mice), ketoconazole did not exhibit any genotoxic potential.

At oral dose levels of 75 mg/kg/day (4.5 times the expected topical human dose in mg/m²), ketoconazole impaired reproductive performance and fertility when administered to male rats (increased abnormal sperm, decreased sperm mobility and decreased pregnancy in mated females).

14 CLINICAL STUDIES

The safety and efficacy of EXTINA[®] Foam were evaluated in a randomized, double-blind, vehicle-controlled study in subjects 12 years and older with mild to severe seborrheic dermatitis. In the study, 427 subjects received EXTINA[®] Foam and 420 subjects received vehicle foam. Subjects applied EXTINA[®] Foam or vehicle foam twice daily for 4 weeks to affected areas on the face, scalp, and/or chest. The overall disease severity in terms of erythema, scaling, and induration was assessed at Baseline and week 4 on a 5-point Investigator's Static Global Assessment (ISGA) scale.

Treatment success was defined as achieving a Week 4 (end of treatment) ISGA score of 0 (clear) or 1 (majority of lesions have individual scores for scaling, erythema, and induration that averages 1 [minimal or faint]) and at least two grades of improvement from baseline. The results are presented in Table 2. The database was not large enough to assess whether there were differences in effects in age, gender, or race subgroups.

Table 2: Efficacy Results

| Number of Subjects | EXTINA [®] Foam N = 427 n (%) | Vehicle Foam N = 420 n (%) |
|--------------------------------------|--|----------------------------------|
| Subjects Achieving Treatment Success | 239 (56%) | 176 (42%) |

16 HOW SUPPLIED/STORAGE AND HANDLING

EXTINA[®] Foam, 2% is supplied in 50 g (NDC 63032-051-50) and 100 g (NDC 63032-051-00) aluminum containers.

Store at controlled room temperature 68°–77°F (20°–25°C). Do not store under refrigerated conditions. **Do not expose containers to heat, and/or store at temperatures above 120°F (49°C).** Do not store in direct sunlight.

Contents are flammable.

Contents under pressure. Do not puncture and/or incinerate container.

Keep out of reach of children.

17 PATIENT COUNSELING INFORMATION

See FDA-approved patient labeling.

17.1 Instructions for Use

- **Avoid fire, flame and/or smoking during and immediately following application.**
- Do not apply EXTINA[®] Foam directly to hands. Dispense onto a cool surface, and apply to the affected areas using the fingertips.

17.2 Local Reactions

- EXTINA[®] Foam may cause skin irritation (application site burning and/or reactions)
- EXTINA[®] Foam may cause contact sensitization.
- As with any topical medication, patients should wash their hands after application.
- Inform a physician if the area of application shows signs of increased irritation and report any signs of adverse reactions.

PATIENT INFORMATION

EXTINA[®] (ex-TEEN-ah) Foam (ketoconazole, 2%)

IMPORTANT: For skin use only. Do not use in the eyes, mouth or vagina.

Read the Patient Information that comes with EXTINA[®] Foam before you start using it and each time you get a refill. There may be new information. This leaflet does not take the place of talking with your doctor about your condition or treatment.

What is EXTINA[®] Foam?

EXTINA[®] Foam is used on the skin (topical) to treat a skin condition called seborrheic dermatitis in patients 12 years and older. Seborrheic dermatitis can cause areas of flaky skin (scales) on the scalp, face, ears, chest or upper back.

EXTINA[®] Foam has not been studied in children less than 12 years old.

What should I tell my doctor before using EXTINA[®] Foam?


For female patients, tell your doctor if you:

- **are pregnant or become pregnant.** It is not known if EXTINA[®] Foam can harm a fetus (unborn baby).
- **breastfeeding.** It is not known if EXTINA[®] Foam passes into breast milk.

How should I use EXTINA[®] Foam?

- **Apply EXTINA[®] Foam exactly as prescribed.** EXTINA[®] Foam is usually applied to the affected skin areas two times a day (once in the morning and once at night) for 4 weeks. Talk to your doctor if your skin does not improve after 4 weeks of treatment with EXTINA[®] Foam.
- **Keep the EXTINA[®] Foam can away from and do not spray it near fire, open flame, or direct heat. EXTINA[®] Foam is flammable. Never throw the EXTINA[®] Foam can into a fire, even if the can is empty.**


Instructions for applying EXTINA[®] Foam

| | |
|---|--|
|  | 1. Hold the can at an upright angle. |
| | 2. Push the button to spray EXTINA [®] Foam directly into the cap of the can or other cool surface. Spray only the amount of EXTINA [®] Foam that you will need to cover your affected skin. Do not spray EXTINA[®] Foam directly onto your affected skin or your hands because the foam will begin to melt right away when it touches your skin. |
| | 3. If your fingers are warm, rinse them in cold water first. Be sure to dry them well before handling the EXTINA [®] Foam. If the EXTINA [®] Foam can seems warm or the foam seems runny, place the can under cool running water for a few minutes. |



4. Using your fingertips, gently massage EXTINA[®] Foam into the affected areas until the foam disappears.





5. If you are treating skin areas with hair such as your scalp, move any hair away so that the foam can be applied to the affected skin.
6. **Do not get EXTINA[®] Foam in your eyes, mouth or vagina. If any EXTINA[®] Foam gets in your eyes, mouth or vagina, rinse areas well with water.**
7. Wash your hands well after applying EXTINA[®] Foam.

What are the possible side effects of EXTINA[®] Foam?

The most common side effects of EXTINA[®] Foam are reaction or burning on treated skin areas. Tell your doctor if you have any reaction on your treated skin such as redness, itching, or a rash. These are not all the side effects of EXTINA[®] Foam. Ask your doctor or pharmacist for more information.

How should I store EXTINA[®] Foam?

- **EXTINA[®] Foam is flammable.**
- **Do not spray EXTINA[®] Foam near fire or direct heat. Never throw the can into a fire, even if the can is empty.**
- **Store the can of EXTINA[®] Foam at room temperature, 68° to 77°F (20°-25°C).** Do not place the EXTINA[®] Foam can in the refrigerator or freezer.
- **Keep the EXTINA[®] Foam can away from all sources of fire and heat. Do not leave the EXTINA[®] Foam can in direct sunlight.**
- **Do not smoke while holding the EXTINA[®] Foam can or while spraying or applying the foam.**
- **Do not pierce or burn the EXTINA[®] Foam can.**
- **Keep EXTINA[®] Foam and all medicines out of the reach of children.**

General information about EXTINA[®] Foam

Medicines are sometimes prescribed for conditions that are not mentioned in Patient Information leaflets. Do not use EXTINA[®] Foam for any other condition for which it was not prescribed. Do not give EXTINA[®] Foam to other people, even if they have the same condition that you have. It may harm them.

This leaflet summarizes the most important information about EXTINA[®] Foam. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about EXTINA[®] Foam that is written for health professionals.

If you have questions about EXTINA[®] Foam you can also call: 1-888-500-DERM (this is a toll-free number) between 6:00 a.m. and 4:00 p.m. Pacific Standard Time, Monday through Friday.

What are the ingredients in EXTINA[®] Foam?

Active ingredients: ketoconazole

Inactive Ingredients: cetyl alcohol, citric acid, ethanol (denatured with tert-butyl alcohol and brucine sulfate) 58%, polysorbate 60, potassium citrate, propylene glycol, purified water, and stearyl alcohol pressurized with a hydrocarbon (propane/butane) propellant.

Rx Only

This Patient Information leaflet has been approved by the U.S. Food and Drug Administration.

The Patient Information leaflet was last revised: November 2008

Manufactured for

Stiefel Laboratories, Inc., Coral Gables, FL 33134 USA

VERSAFOAM-HF, VERSAFOAM-HF and Design, EXTINA, and V Design, STIEFEL, and STIEFEL and Design are registered trademarks of Stiefel Laboratories, Inc.

U.S. Patent Pending

©2008 Stiefel Laboratories, Inc.

302418

AW-0854

Principal Display Panel

NDC 63032-051-50

extina[®]

(ketoconazole) Foam, 2%

50 g R_x only

FOR TOPICAL USE ONLY

NOT FOR OPHTHALMIC, ORAL OR INTRAVAGINAL USE.

CFC FREE

Description: Extina (ketoconazole) Foam, 2% contains 20 mg ketoconazole, USP per gram in a thermolabile hydroethanolic foam vehicle consisting of cetyl alcohol NF, citric acid USP, ethanol (denatured with *tert*-butyl alcohol and brucine sulfate) 58%, polysorbate 60 NF, potassium citrate USP, propylene glycol USP, purified water USP, stearyl alcohol NF pressurized with a hydrocarbon (propane/butane) propellant.

Hold the can at an upright angle and then press firmly to dispense.

Dosage: Use only as prescribed by your physician. See package insert for full prescribing information.

Warning: FLAMMABLE. AVOID FIRE, FLAME, OR SMOKING DURING AND IMMEDIATELY FOLLOWING APPLICATION.

Contents under pressure. Do not puncture and/or incinerate container. Do not expose containers to heat, and/or store at temperatures above 120°F (49°C). Do not store in direct sunlight.

Avoid contact with eyes. Keep out of reach of children.

Do not store in the refrigerator.

Store at controlled room temperature, 68° – 77°F (20° - 25°C).

Manufactured for

Stiefel Laboratories, Inc.

Coral Gables, FL 33134

USA

For additional information:

1-888-500-DERM or visit

www.extina.com

DELIVERED IN VersaFoam-HFTM

HYDROETHANOLIC FORMULATION

AW No: AW-0591-r3

P/N: 128573-0607

©2007, Stiefel Laboratories, Inc.

VersaFoam-HF is a trademark, and Extina, the V logo and Stiefel are registered trademarks, owned by Stiefel Laboratories, Inc.

U.S. Patent Pending

Recyclable Aluminum Container

DESCRIPTION

Extina® (ketconazole) Foam, 2% contains 20 mg ketconazole, USP per gram in a thermally stable hydroalcoholic foam vehicle consisting of cetyl alcohol NF, cetyl estol USP, ethanol (denatured with tert-butyl alcohol and benzoin suffragin) 50%, polyacrylate 10 NF, potassium chloride USP, propylene glycol USP, purified water USP, stearyl alcohol NF presaturated with a hydrocarbon (gasoline/butane) propellant.

Manufactured for
Stiefel Laboratories, Inc.
Coral Gables, FL 33134
USA

For additional information:
1-800-500-DERM or visit
www.extina.com

ADV No.: ADV-0091-03

P/N: 128573-0503

© 2007 Stiefel Laboratories, Inc.

Extina® is a trademark, and Extina, the V logo, and Stiefel are registered trademarks owned by Stiefel Laboratories, Inc.

U.S. Patent Pending



NDC 63032-051-50
50 g Rx only
FOR TOPICAL USE ONLY

extina®
(ketconazole) Foam, 2%

MANUFACTURED IN



VerelFoam®
Pharmaceuticals, Inc.

**NOT FOR OPHTHALMIC,
ORAL, OR INTRAPERITONEAL USE.**

Hold the can at an
upright angle and
then press firmly to
dispense.



Warnings: Use only as prescribed by
your physician. See package insert
for full prescribing information.

**Warnings: FLAMMABLE. AVOID
FIRE, FLAME, OR EXCESSIVE
HEATING AND IMMEDIATELY
FOLLOWING APPLICATION.**

Contents under pressure. Do not
puncture and/or incinerate
containers. Do not expose
containers to heat, sunlight or
other temperatures above 120°F
(49°C). Do not store in direct
sunlight.

Avoid contact with eyes. Keep out
of reach of children.

Do not store in the refrigerator.
Store at controlled room
temperature, 68°-77°F (20°-25°C).



3 63032 05150 2

Revised: 10/2009

Distributed by: Stiefel Laboratories Inc